

Our undiluted antisera are suitable for immunohistochemical use, for application to radioimmunoassay (RIA) and other non-isotopic immunoassay systems. While these antisera are characterized in terms of their specificity and sensitivity using enzyme immunoassays, they are not controlled to the rigorous specifications of prediluted antisera. These can be adjusted to optimum concentrations for RIA or other immunoassay.

**14216-v SPAI-1 (Porcine) Antiserum**  
**Sodium Potassium ATPase Inhibitor-1 (Porcine) Antiserum**  
**Na<sup>+</sup>, K<sup>+</sup>-ATPase Inhibitor-1 (Porcine) Antiserum**  
(Rabbit)

Antiserum 50 µl/vial  
Lyophilized from 0.001 M Phosphate Buffer (pH 7.0)

Lot No. 901-400318

Immunogen : SPAI-1 (Porcine) - BSA (BSA : Bovine Serum Albumin)

Specificity :

SPAI-1 (Porcine)	100%
SPAI-1 (Porcine, 1-14)	0%
SPAI-1 (Porcine, 25-35)	0%
Ouabain	0%

No cross-reactions against the following peptides are observed at much higher doses (100-200 pmol/ml).  
Calcitonin (Human), CRF (Human), Endothelin-1 (Human), BNP-26 (Porcine), and Secretin (Human)


Sensitivity : IC<sub>50</sub> = 79 pg/tube, (0.14 pmol/ml)

The standard curve is obtained by enzyme immunoassay using β-D-galactosidase as a label. A brief diagram of the assay system is shown below.

Storage and Stability :

This lyophilized antiserum is stable for one month at room temperature. For long time storage, we recommend storing it in a deep freezer. After reconstitution with your buffer, the solution can be kept at 4° for continuous use with a preservative. For extended storage of the solution, it should be frozen at -20°.

**Standard Curve of SPAI-1 (Porcine)**



**Assay System**

Diluted antiserum	0.1 ml
Standard solution	0.1 ml
Assay buffer	0.1 ml

↓

Preincubation at 4° for 5-6 hr.

↓

Delayed addition of β-D-galactosidase  
labeled antigens 0.1 ml

↓

1st Incubation at 4° for 15 hr.

↓

Addition of immunobead 0.1 ml

↓

2nd Incubation at room temperature for 3 hr.

↓

Centrifugation at 4° for 5 min.

↓

Decantation and rinse (twice)

↓

Determination of enzyme activity in precipitate

↓

Calculation (Bx/Bo x 100)

**Caution: This product is sold for research purposes only and not for use in humans.**